ProSci LLC
12170 Flint Place
Poway, CA 92064, USA
ProSciAntibodies.com

P: +1 (888) 513-9525
Local: +1 (858) 513-2638
Fax: +1 (858) 513-2692

CustomerCare@ProSciAntibodies.com
ITSupport@ProSciAntibodies.com

TIMP2 Rabbit Recombinant Antibody (82-272)


CATALOG NUMBER: 82-272

Specifications

Host Species	Rabbit
Species Reactivity	African Green Monkey, Human, Mouse, Rat
Immunogen	Synthetic peptide. This information is considered to be commercially sensitive.
Conjugate	Unconjugated
Tested Applications	ELISA, IF/ICC, WB
User Note	Optimal dilutions for each application to be determined by the researcher.
Predicted Molecular Weight	24 kDa


Properties

Purification	Affinity purification
Clonality	Monoclonal
Isotype	IgG
Physical State	Liquid
Buffer	Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.
Concentration	Batch dependent
Storage Conditions	Store at -20°C. Avoid freeze / thaw cycles.




ARC51192-01_WB_01_82-272

Western blot analysis of various lysates using TIMP2 Rabbit mAb at 1:1000 dilution incubated overnight at 4°C. | Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1:10000 dilution. | ...




ARC51192-01_IF_01_82-272

Confocal imaging of SH-SY5Y cells using TIMP2 Rabbit mAb followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.



ARC51192-01_IF_02_82-272

Confocal imaging of SW480 cells using TIMP2 Rabbit mAb followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.



ARC51192-01_IF_03_82-272

Confocal imaging of NIH/3T3 cells using TIMP2 Rabbit mAb followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (Red). The cells were counterstained with α-Tubulin Mouse mAb followed ...

Disclaimer

Disclaimer	This product is for research use only.

For research use only. For additional information, visit ProSci's Terms and Conditions Page.